This listing of claims will replace all prior versions, and listings, of the claims in the application.

Listing of Claims

1. (currently amended) An ancient defense polymer having selective antimicrobial activity, and comprising:

A) one or more discrete hydrophobic segments, wherein said hydrophobic segment comprises:

A1. polymerized hydrophobic chain growth monomers;

A2. polymerized step-growth monomers; or

A3. hydrophobic (di)functional oligomers or polymers; and

B) one or more hydrophilic segments containing cationic functionality wherein said hydrophilic segment comprises:

B1. polymerized cationic chain growth monomers;

B2. a polymer made from a mixture of cationic chain growth monomers and (i) uncharged monomers that are hydrophilic or (ii) hydrophobic monomers; or

B3. cationic (di)functional oligomers or polymers;

wherein the ancient defense polymer has a grafted chain architecture, comprising a main chain and chains grafted onto the main chain, wherein the main chain comprises at least one of said hydrophilic segments and the graft chains comprise at least one of said hydrophobic segments, and

wherein the main chain is not cellulose, and

wherein the ancient defense polymer has selective antimicrobial activity, has low aqueous solubility, and high molecular weight.

- 2. (cancelled)
- 3. (cancelled)

Appln. No. 10/552,714

Response dated: March 31, 2010

Reply to Office Action of January 15, 2010

4. (previously presented) The ancient defense polymer of claim 1, wherein said hydrophobic

segment comprises polymerized hydrophobic alkyl methacrylates, aryl methacrylates, alkyl

methacrylamides, or aryl methacrylamides.

5. (previously presented) The ancient defense polymer of claim 1, wherein said hydrophilic

segment comprises polymerized methacrylates and/or methacrylamides.

6. (previously presented) The ancient defense polymer according to claim 1 comprising a

copolymer of 3-aminopropyl methacrylamide (AMA,) and poly(propylene

oxide)monomethacrylate (PPO-ME).

7. (previously presented) The ancient defense polymer according to claim 6 wherein AMA is

present in an amount of from 5 to 50 mol%.

8. (previously presented) The ancient defense polymer according to claim 6, wherein AMA is

present in an amount of about 10 mol%.

9. (original) The ancient defense polymer according to claim 1, comprising a terpolymer of 3-

aminopropyl methacrylamide (AMA), poly(propylene oxide)monomethacrylate (PPO-ME), and

methyl methacrylate.

10. (original) The ancient defense polymer according to claim 1, comprising a terpolymer

of 3-aminopropyl methacrylamide (AMA), poly(propylene oxide)monomethacrylate (PPO-ME),

and n-butyl methacrylate (BMA).

11. (cancelled)

12. (cancelled)

13. (previously presented) The ancient defense polymer of claim 1, wherein the hydrophilic

and/or the hydrophobic segments are grafted either directly or indirectly to the main chain, and

the main chain is a derivatizable polymer.

Page 3 of 8

Appln. No. 10/552,714

Response dated: March 31, 2010

Reply to Office Action of January 15, 2010

14. (previously presented) The ancient defense polymer according to claim 13, wherein at least one of said hydrophobic segments or said hydrophilic segments is grafted to the derivatizable polymer by a spacer group.

- 15. (previously presented) The ancient defense polymer according to claim 13, wherein said hydrophobic segment and said hydrophilic segment are grafted to the derivatizable polymer by a spacer group.
- 16. (cancelled)
- 17. (original) The ancient defense polymer according to claim 13, wherein said hydrophobic segment is grafted onto said hydrophilic segment.
- 18. (previously presented) The ancient defense polymer of claim 13, wherein the derivatizable polymer comprises polymerized chain growth monomers containing reactive functional groups.
- 19. (previously presented) The ancient defense polymer of claim 18, wherein said functional groups are one or more of the groups selected from the group consisting of hydroxyl, carboxylic acid, amine, vinyl, acid chloride, and isocyanate.
- 20. (currently amended) An apparatus comprising an ancient defense polymer having selective antimicrobial activity and comprising:
- A) one or more discrete hydrophobic segments, wherein said hydrophobic segment comprises:
 - A1. polymerized hydrophobic chain growth monomers;
 - A2. polymerized step-growth monomers; or
 - A3. hydrophobic (di)functional oligomers or polymers; and
- B) one or more hydrophilic segments containing cationic functionality wherein said hydrophilic segment comprises:
 - B1. polymerized cationic chain growth monomers;
 - B2. a polymer made from a mixture of cationic chain growth monomers and (i) uncharged monomers that are hydrophilic or (ii) hydrophobic monomers; or

B3. cationic (di)functional oligomers or polymers;

wherein said ancient defense polymer is bound in or attached to a surface of said apparatus to impart antimicrobial activity to said apparatus,

wherein the main chain comprises at least one of said hydrophilic segments and the graft chains comprise at least one of said hydrophobic segments, and

wherein the main chain is not cellulose, and

wherein the ancient defense polymer has selective antimicrobial activity, has low aqueous solubility, and high molecular weight.

- 21. (previously presented) The apparatus of claim 20, wherein said apparatus is selected from the group consisting of an implant, a catheter, a replacement valve, a wound dressing, a medical device, and a stent.
- 22. (previously presented) The ancient defense polymer of claim 10, made from 1-15 mol% BMA, 5-49 mol% AMA, and 50-90 mol% PPO-Me.
- 23. (previously presented) The apparatus of claim 21, wherein the ancient defense polymer has a grafted chain architecture, comprising a main chain and chains grafted onto the main chain.
- 24. (currently amended) An apparatus consisting of, or having a portion consisting of, an ancient defense polymer having selective antimicrobial activity and comprising:
- A) one or more discrete hydrophobic segments, wherein said hydrophobic segment comprises:
 - A1. polymerized hydrophobic chain growth monomers;
 - A2. polymerized step-growth monomers; or
 - A3. hydrophobic (di)functional oligomers or polymers; and
- B) one or more hydrophilic segments containing cationic functionality wherein said hydrophilic segment comprises:
 - B1. polymerized cationic chain growth monomers;
 - B2. a polymer made from a mixture of cationic chain growth monomers and (i) uncharged monomers that are hydrophilic or (ii) hydrophobic monomers; or
 - B3. cationic (di)functional oligomers or polymers;

wherein said ancient defense polymer imparts antimicrobial activity to said apparatus,

wherein the main chain comprises at least one of said hydrophilic segments and the

graft chains comprise at least one of said hydrophobic segments, and

wherein the main chain is not cellulose, and

wherein the ancient defense polymer has selective antimicrobial activity, has low aqueous solubility, and high molecular weight.

- 25. (previously presented) The apparatus of claim 24, selected from the group consisting of surgical devices, sterile draping and dressings, clothing, food packaging, agricultural processing and bioreactor parts.
- 26. (previously presented) The apparatus of claim 25, wherein the ancient defense polymer has a grafted chain architecture, comprising a main chain and chains grafted onto the main chain.
- 27. (cancelled)
- 28. (cancelled)
- 29. (cancelled)
- 30. (cancelled)
- 31. (new) The ancient defense polymer of claim 1, having a molecular weight of at least 143,970.